# **Mycterothrips**

## Generic diagnosis

Female macropterous. Head often slightly narrowed to base; maxillary palps 3-segmented; eyes with five usually weakly pigmented facets; mouth-cone sometimes long or even extending to mesosternum; ocellar setae I usually present, setae I and II sometimes absent, setae III long within ocellar triangle; five pairs of postocular setae. Antennae 8-segmented (rarely 7segmented); segment I with paired dorso-apical setae, III and IV with forked sense-cones, III-VI with some microtrichial rows on both surfaces. Pronotum with two pairs of long posteroangular setae; two pairs of posteromarginal setae. Mesonotum with median pair of setae near or far from posterior margin; campaniform sensilla absent. Metanotum longitudinally or irregularly reticulate; median pair of setae usually at or near anterior margin; campaniform sensilla absent. Fore wing first vein with long gap in setal row, about seven or eight basal and two distal setae, second vein with evenly spaced setal row, posterior fringe cilia wavy; clavus with about five veinal and one discal setae. Prosternal ferna connected weakly at middle; basantra membranous, without setae; prospinasternum broad and transverse. Mesosternopleural sutures weak and not reaching anterior margin; endofurca with spinula. Metathoracic endofurca usually with well-developed spinula. Tarsi 2-segmented. Tergites without ctenidia or craspeda; II-VIII laterally, also laterotergites, with rows of microtrichia, or with scattered dentate microtrichia; tergite VIII with comb long, fine and complete; tergite IX with two pairs of campaniform sensilla, X with or without median split. Sternites without craspeda, often with discal setae; III-VII usually with three pairs of posteromarginal setae, II with two pairs; laterotergites without discal setae.

Male similar to female, but rarely wings absent, antennae in some species 6or 7-segmented with segment VI much longer than in female with long





consociatus head & thorax consociatus antenna



consociatus dorso-apical setae on antennal



consociatus tergites VII-VIII





consociatus meso & meta furca glycines head & pronotum glycines tergites VII-IX



consociatus fore wing

lateral setae; tergites with fewer microtrichia; sternites with or without discal setae, without pore plates.

#### **Relationship data**

Thripidae sub-family Thripinae: this is a diverse group involving more than 230 genera. This genus is one of only 16 genera from China in which species have a pair of dorso-apical setae on the first antennal segment. However, considering the diversity amongst these genera, including for example genera related to *Trichromothrips* and genera related to *Odontothrips*, this character state is not necessarily indicative of monophyly.

#### **Biological data**

The species apparently all feed and breed on leaves, although adults can disperse widely onto many plants.

#### **Distribution data**

This genus has an extensive distribution around the world. At one time considered essentially Holarctic, many species have now been described extending south across tropical Asia and with two known from Australia and two from Africa.

### Nomenclatural data

Mycterothrips Trybom, 1910: 158. Type species Mycterothrips laticauda Trybom, 1910, by monotypy.

There are 36 species listed in this genus (ThripsWik, 2018), and the following eight are recorded from China:

araliae (Takahashi, 1936: 434). (*Taeniothrips*) auratus Wang, 1999: 232. caudibrunneus Wang, 1999: 234. consociatus (Targioni-Tozzetti, 1887: 425). (*Thrips: Euthrips*) glycines (Okamoto, 1911: 221). (*Euthrips*) gongshanensis Li, Li & Zhang, 2017: 372. nilgiriensis (Ananthakrishnan, 1960: 37). (*Rhopalandrothrips*) ricini (Shumsher, 1946: 176). (*Taeniothrips: Rhopalandrothrips*) setiventris (Bagnall, 1918: 61). (*Physothrips*) yelangi Li, Li & Zhang, 2017: 374.

#### References

Li YJ, Li ZY & Zhang HR (2017) Two new species of *Mycterothrips* (Thysanoptera: Thripidae) from Southwestern China. *Zootaxa* **4353** (2): 371–377.

Masumoto M & Okajima S (2006) A revision of and key to the world species of *Mycterothrips* Trybom (Thysanoptera, Thripidae). *Zootaxa* **1261**: 1–90.

ThripsWiki (2018). ThripsWiki - providing information on the World's thrips. < http://thrips.info/wiki/Main\_Page>

Copyright © 2018. All rights reserved.